

## Anti-oxidative effects of pomegranate juice (PJ) consumption by diabetic patients on serum and on macrophages

Mira Rosenblat<sup>a</sup>, Tony Hayek<sup>a,b</sup>, Michael Aviram<sup>a,\*</sup>

<sup>a</sup> *The Lipid Research Laboratory, Technion Faculty of Medicine, The Rappaport Family Institute for Research in the Medical Sciences, Rambam Medical Center, 31096 Haifa, Israel*

<sup>b</sup> *Internal Medicine Department E, Rambam Medical Center, Haifa, Israel*

Received 14 July 2005; received in revised form 18 July 2005; accepted 10 September 2005

Available online 13 October 2005

### Abstract

Diabetes is associated with increased oxidative stress and atherosclerosis development. In the present study, we investigated the effects of pomegranate juice (PJ; which contains sugars and potent anti-oxidants) consumption by diabetic patients on blood diabetic parameters, and on oxidative stress in their serum and macrophages. Ten healthy subjects (controls) and 10 non-insulin dependent diabetes mellitus (NIDDM) patients who consumed PJ (50 ml per day for 3 months) participated in the study. In the patients versus controls serum levels of lipid peroxides and thiobarbituric acid reactive substances (TBARS) were both increased, by 350% and 51%, respectively, whereas serum SH groups content and paraoxonase 1 (PON1) activity, were both decreased (by 23%). PJ consumption did not affect serum glucose, cholesterol and triglyceride levels, but it resulted in a significant reduction in serum lipid peroxides and TBARS levels by 56% and 28%, whereas serum SH groups and PON1 activity significantly increased by 12% and 24%, respectively. In the patients versus controls monocytes-derived macrophages (HMDM), we observed increased level of cellular peroxides (by 36%), and decreased glutathione content (by 64%). PJ consumption significantly reduced cellular peroxides (by 71%), and increased glutathione levels (by 141%) in the patients' HMDM. The patients' versus control HMDM took up oxidized LDL (Ox-LDL) at enhanced rate (by 37%) and PJ consumption significantly decreased the extent of Ox-LDL cellular uptake (by 39%). We thus conclude that PJ consumption by diabetic patients did not worsen the diabetic parameters, but rather resulted in anti-oxidative effects on serum and macrophages, which could contribute to attenuation of atherosclerosis development in these patients.

© 2005 Elsevier Ireland Ltd. All rights reserved.

**Keywords:** Pomegranate juice; Diabetes; Oxidative stress; Macrophages; Atherosclerosis

### 1. Introduction

Diabetes mellitus is increasing worldwide, resulting from the interaction of obesity, inflammation and hyperglycemia. Both type I and type II diabetes are powerful and independent risk factors for coronary artery disease, stroke and peripheral arterial disease [1,2], and atherosclerosis accounts for 80% of all deaths among diabetic patients. Prolonged exposure to hyperglycemia is now recognized as a major risk factor in the pathogenesis of atherosclerosis in diabetes [3]. Animal and human studies elucidated three major mechanisms

for the pathological alterations observed in diabetic vasculature, i.e. non-enzymatic glycosylation of proteins and lipids which can interfere with their normal function, cellular protein kinase C (PKC) activation and oxidative stress [3,4]. Diabetic patients may be highly prone to oxidative stress because hyperglycemia depletes natural anti-oxidants and facilitates the production of free radicals [5,6]. Thus, anti-oxidants treatment in diabetes could be beneficial [7]. Indeed, it was shown that alpha-tocopherol or red wine supplementation to diabetic patients significantly reduced serum oxidative stress [8,9]. Furthermore, tea catechins were able to protect diabetic erythrocytes from *tert*-butyl hydroperoxide-induced oxidative stress [10]. Pomegranate juice (PJ) possesses impressive anti-oxidative properties due to its polyphenolics, tannins and

\* Corresponding author. Tel.: +972 4 8542970; fax: +972 4 8542130.

E-mail address: aviram@tx.technion.ac.il (M. Aviram).

anthocyanins [11,12]. We have previously demonstrated that consumption of PJ by humans for a period of 1 year significantly reduced the oxidation of both LDL and HDL [13]. Furthermore, in patients with carotid artery stenosis that consumed PJ for 3 years, we demonstrated reduced oxidative stress in their blood, and a decreased atherosclerotic lesion size [14].

Macrophages play a major role in the early stages of atherogenesis [15]. Recent studies that were performed in control subjects or in diabetic patients' monocytes–macrophages demonstrated that high glucose levels can lead to macrophage foam cell formation by several mechanisms including: increased cholesterol synthesis [16], altered expression and secretion of lipoprotein lipase [17], monocytes PKC activation [18] and up-regulation of an oxidized LDL (LOX-1) receptor [19], or scavenger receptors [20]. Recently, we have shown increased oxidative stress and increased uptake of Ox-LDL also in peritoneal macrophages from streptozotocin-induced diabetic mice, as well as in vitro, in cells that were incubated with high glucose levels [21].

The diabetic patients avoid sugar-containing juices, such as grape juice, which can worsen their diabetic conditions. In the present study, we questioned whether PJ (which contains 10% sugars and potent polyphenols anti-oxidants) consumption by diabetic patients also worsen diabetes and its oxidative complications.

## 2. Methods

### 2.1. Subjects

Ten male healthy subjects (controls) and 10 male non-insulin dependent diabetes mellitus (NIDDM) patients (age 35–71 years old, mean age  $50 \pm 10$ ) participated in the study. The controls were non-smokers, with no diabetes (glucose levels below 100 mg% and hemoglobin A1c levels were in the range of 4.8–6.2%), hypertension or coronary artery disease, and they did not take any medications. The diabetes mellitus duration in the patients was 4–10 years, glucose levels above 160 mg%, hemoglobin A1c in the range of 7.5–11.3%. All the patients had no ischemic heart disease, no hypercholesterolemia and were no smokers, but 50% of the patients were hypertriglyceridemic with serum triglyceride levels (300–790 mg%). Eighty percent of the patients were treated with Glucophage (Metformin) and 50% with Gluben (Glybenclamid). Two of the patients were hypertensive and were treated with ACE inhibitors. The patients consumed 50 ml of pomegranate juice per day (which contain 1.5 mmol of total polyphenols) for a period of 3 months. Blood was collected from controls and from the diabetic patients before and after PJ consumption for biochemical parameters analysis. Blood was also collected from two of the controls and from three diabetic patients before and after PJ consumption for preparation of monocytes-derived macrophages.

### 2.2. Reagents

2',7'-Dichlorofluorescein diacetate (DCFH) was purchased from Sigma (St. Louis, MO, USA). FITC-conjugated antibody was purchased from Serotec IQ Products (Zerinkepark, The Netherlands). PBS, DMEM, RPMI-1640 medium, FCS (heat-inactivated at 56 °C for 30 min), penicillin, streptomycin, nystatin, L-glutamine and sodium pyruvate were purchased from Biological Industries (Beth Haemek, Israel).

### 2.3. Pomegranate processing

Pomegranates were picked by hand, washed and stored in tanks. The fruits were crushed and squeezed. The juice was filtered, pasteurized, concentrated and stored at  $-18$  °C. Each day along the study period, the concentrated PJ was diluted 1:5 (v/v) with water in order to obtain a single strength PJ. The anti-oxidant composition of the juice includes: 1979 mg/l of tannins (1561 mg/l of punicalagin and 417 mg/l of hydrolysable tannins), 384 mg/l of anthocyanins (delphinidin 3,5-diglucoside, cyanidin 3,5-diglucoside, delphinidin-3-glucoside, cyanidin 3-glucoside and pelargonidine 3-glucoside) and 121 mg/l of ellagic acids derivatives. The juice contained also 3 mg of Vitamin C per 100 ml of PJ. For the extraction of PJ polyphenols fraction C18 sorbent column was used (Varian HF Bondesil C18 resin sorbent). Total polyphenols were eluted from the column with 1% acidified (food-grade acetic acid) ethanol.

### 2.4. Serum paraoxonase 1 activity

PON1 arylesterase activity towards phenyl acetate was determined as previously described [22].

### 2.5. Serum lipids peroxidation

Serum lipid peroxidation was measured before and after 3 months of PJ consumption. Serum samples were diluted  $\times 4$  with PBS, and were incubated without or with 100 mM of 2,2'-azobis, 2-amidinopropane hydrochloride (AAPH, Wako, Japan) for 2 h at 37 °C [23]. The extent of lipid peroxidation was measured by the thiobarbituric acid reactive substances (TBARS) assay [24] and by the lipid peroxides assay [25].

### 2.6. Total thiols (SH groups) in serum

The assay procedure determines the amount of protein bound SH groups, as well as glutathione [26]. An aliquot of 50  $\mu$ l serum was mixed with 1 ml of Tris–EDTA buffer, and the absorbance at 412 nm was measured. To this was added 20  $\mu$ l of 10 mM DTNB, and after 15 min incubation at room temperature the absorbance was measured, together with a DTNB blank. Total SH groups are calculated as described before [26].

### 2.7. Human monocytes-derived macrophages (HMDM)

HMDM were separated from the blood [17] and plated at  $10^6$ /ml in RPMI medium with 10% FCS. After 2 h of incubation at 37 °C, non-adherent cells were removed, and RPMI with 10% autologous serum was added. Macrophages were analyzed 8 days after plating.

### 2.8. Detection of intracellular oxidative stress by the DCFH assay

Intracellular oxidative stress was assayed through the oxidation of DCFH-DA [27], and monitored by flow cytometry [28]. For flow-cytometric assay of DCFH-DA oxidation, cells were washed ( $\times 1$ ) with PBS and incubated with 10  $\mu$ M DCFH-DA, in medium for 30 min at 37 °C. Adherent cells were detached by gentle scraping, and all cells were washed ( $\times 2$ ) with PBS. Measurements of cellular fluorescence determined by FACS were done at 510–540 nm after excitation of the cells at 488 nm with an argon ion laser. Ten thousand events were registered for each experiment. Cellular fluorescence was quantitated by mean fluorescence intensity (MFI).

### 2.9. Macrophage reduced glutathione content

All the preparation steps were carried out on ice. The cells from triplicate dishes ( $1 \times 10^6$  per dish) were washed, scraped from the dish and sonicated in an ultrasonic processor ( $3 \times 20$  s at 80 W). The amount of protein was measured by the Lowry method [29] and reduced glutathione content by the DTNB-GSSG reductase recycling assay [30].

### 2.10. Oxidized LDL (Ox-LDL) uptake by macrophages

LDL was separated from plasma of normal healthy volunteers by discontinuous density-gradient ultracentrifugation [31] and dialyzed against saline with EDTA (1 mM). LDL protein concentration was determined by the Lowry method [29]. Before oxidation, LDL was diluted in PBS to 1 mg/ml and dialyzed overnight against PBS at 4 °C to remove the EDTA. Oxidation of LDL was carried out at 37 °C under air in a shaking water bath. LDL (1 mg/ml) was incubated for 18 h at 37 °C with freshly prepared  $\text{CuSO}_4$  (5  $\mu$ M, Sigma).

Oxidation was terminated by refrigeration at 4 °C. The extent of LDL oxidation was determined by the TBARS assay [24]. Ox-LDL was conjugated to fluoroisothiocyanate (FITC) for cellular uptake studies. HMDM were incubated at 37 °C for 3 h with FITC-conjugated Ox-LDL at a concentration of 20  $\mu$ g of protein/ml. The uptake of the lipoproteins was determined by flow cytometry. Measurements of cellular fluorescence determined by FACS were done at 510–540 nm after excitation of the cells at 488 nm with an argon ion laser. Ten thousand events were registered for each experiment. Cellular fluorescence was quantitated by mean fluorescence intensity.

### 2.11. Statistical analysis

Statistical analysis was performed using the Student paired *t*-test when comparing the mean of two groups. ANOVA was used when more than two groups were compared and results are given as mean  $\pm$  S.E.M.

## 3. Results

### 3.1. Effect of PJ consumption by diabetic patients on serum biochemical parameters

Serum total cholesterol and LDL cholesterol levels in the patients were similar to that of the controls. In contrast, serum triglyceride levels were significantly higher by 2.8-fold in the patients versus controls, whereas HDL-cholesterol levels were significantly decreased by 28% (Table 1). PJ consumption by the patients did not affect these parameters (Table 1). As PJ contains sugars we first questioned the effect of PJ consumption by the patients on serum diabetic parameters: glucose, hemoglobin (Hb) A1c, insulin and C-peptide (a cleavage product of proinsulin). Blood Hb A1c levels were significantly increased in the diabetic patients versus controls by 59% ( $8.9 \pm 0.5\%$  versus  $5.6 \pm 0.2\%$ ), whereas insulin and serum C-peptide levels were only slightly different in patients versus controls ( $82 \pm 9$  pmol/l for the patient's insulin levels versus  $103 \pm 8$  pmol/l for the control's insulin levels, and  $862 \pm 119$  pmol/l for the patient's C-peptide versus  $770 \pm 88$  pmol/l for the control's C-peptide

Table 1  
Serum lipid profile in healthy subjects (controls) and in diabetic patients before and after pomegranate juice (PJ) consumption

	Controls	Diabetic patients before PJ	Diabetic patients after PJ
Triglyceride (mg%)	115 $\pm$ 25	327 $\pm$ 79*	302 $\pm$ 64
Total cholesterol (mg%)	190 $\pm$ 9	203 $\pm$ 13	193 $\pm$ 10
LDL-cholesterol (mg%)	111 $\pm$ 10	112 $\pm$ 8	110 $\pm$ 8
HDL-cholesterol (mg%)	56 $\pm$ 3	39 $\pm$ 3	41 $\pm$ 3*

NIDDM patients ( $n = 10$ ) were compared to healthy subjected (controls,  $n = 10$ ). The patients consumed pomegranate juice (PJ; 50 ml per day for 3 months). Serum samples were collected from the controls and from the patients before and after PJ consumption. The lipid profile in the serum samples was then determined. Results are given as mean  $\pm$  S.E.M. ( $n = 10$  in each group).

\*  $p < 0.01$  vs. controls.




Fig. 1. The effect of PJ consumption by diabetic patients on diabetic parameters in serum. Ten NIDDM patients consumed PJ (50 ml per day for 3 months). Blood samples were collected from the patients before PJ consumption and after 3 months of PJ consumption. Serum fasting glucose (A), hemoglobin A1c (B), insulin (C) and C-peptide (D) levels were determined. Results are given as the individual results, and also as the mean of the whole group, \* $p < 0.01$  vs. before PJ consumption.

levels, respectively). PJ consumption by the patients resulted in a non-significant reduction, by 8%, in the serum glucose levels, with no significant effect on HbA1c levels (Fig. 1A and B). Similarly, a non-significant reduction in serum insulin levels (by 9%) was noted (Fig. 1C), whereas serum C-peptide levels were significantly lower, by 23%, in patients after PJ consumption versus before (Fig. 1D). These results indicate that, in spite of the presence of sugars in consumed PJ, serum diabetic parameters were not worsen, but even improved (Fig. 1).

### 3.2. Effect of PJ consumption by diabetic patients on serum oxidative status

Diabetes is known to be accompanied by increased oxidative stress [4–7]. Indeed, in our patients' serum samples we observed significant high levels of lipid peroxides and TBARS by 350% and 51%, respectively, versus the controls (Fig. 2A and B). In addition serum sulfhydryl groups content (which is another marker for oxidative stress) was significantly reduced by 21% in the patients versus controls (Fig. 2C). Paraonase 1 is an HDL-associated lactonase/esterase which was shown to protect lipids in lipoproteins and cells from oxidation, by its ability to hydrolyze

specific oxidized lipids [32]. PON1 arylesterase activity in the patients' serum was significantly lower, by 23%, versus the controls (Fig. 2D). PJ consumption by the patients significantly reduced serum oxidative stress. The lipid peroxides and TBARS levels were decreased by 56% and 28%, respectively, as compared to the levels observed in the patients' serum before PJ consumption (Fig. 1A and B). In parallel, serum total sulfhydryl groups content and PON1 arylesterase activity, significantly increased by 12% and 24%, respectively (Fig. 2C and D).

### 3.3. Direct effect of PJ on serum oxidative stress: in vitro study

Diabetic patients' serum samples (obtained from the patients before PJ consumption) were incubated for 1 h at room temperature (25 °C) with no addition (control) or with PJ (20  $\mu$ M or 40  $\mu$ M of total PJ polyphenols). Then, the amount of TBARS in the basal state, as well as after AAPH-induced lipid peroxidation, was measured. PJ significantly decreased the basal amount of TBARS in the patients' serum by 17% or 24%, on using 20  $\mu$ M or 40  $\mu$ M of total polyphenols, respectively (Fig. 3A). Furthermore, the susceptibility of the patients' serum to AAPH-induced




Fig. 2. The effect of PJ consumption by diabetic patients on serum oxidative status. Ten NIDDM patients were compared to 10 healthy subjects (controls). The patients consumed PJ (50 ml per day for 3 months). Blood samples were collected from the controls and from the patients before and after PJ consumption. The oxidative status of the serum samples was determined by the lipid peroxides (A) and TBARS (B) assays. Total serum thiols (SH) groups levels (C) and paraoxonase 1 arylesterase activity (D) were measured as described under Section 2. Results are given as mean  $\pm$  S.E.M., \* $p < 0.01$  vs. controls; # $p < 0.01$  after PJ consumption vs. before PJ consumption.

oxidation was also substantially decreased, by 48% and 73%, respectively (Fig. 3B). Upon adding increasing concentrations of PJ total polyphenols (0–40  $\mu$ M) to diabetic patients' serum samples, PON1 arylesterase activity significantly increased in a PJ dose-dependent manner, by up to 25% (Fig. 3C).

#### 3.4. Effect of PJ consumption by diabetic patients on cellular oxidative status in their monocytes-derived macrophages

As diabetic patients are prone to develop accelerated atherosclerosis [1–3], and as macrophages play a major role in the early stages of atherogenesis [15,32], we next studied the oxidative status of the patients' HMDM versus controls, and the effect of PJ consumption by the diabetic patients. The level of total cellular peroxides, as measured by the DCFH assay, was significantly higher by 36% in the patients' HMDM versus controls' HMDM (Fig. 4A). PJ consumption significantly reduced the cellular lipid peroxides content by 71% (to levels which are even lower than those observed in the controls' HMDM), in comparison to the levels observed in the patients' HMDM before PJ consumption (Fig. 4A).

Reduced glutathione (GSH) is a major cellular anti-oxidant against oxidative stress [33]. In the patients' HMDM, reduced glutathione content was markedly lower, by 64%, versus the amount found in controls' HMDM, and PJ consumption by the patients resulted in elevation in the HMDM glutathione levels by 141%, almost the level observed in control HMDM (Fig. 4B). These results indicate that in diabetic patients' macrophages, and not only the serum, are under increased oxidative stress. PJ consumption by diabetic patients demonstrated beneficial effects by reducing cellular oxidative stress in the patients' macrophages.

Oxidized LDL uptake by macrophages can lead to macrophage cholesterol accumulation and foam cell formation [32], and "oxidized macrophages" were shown to take up Ox-LDL at enhanced rate [34]. Thus, we next compared the extent of Ox-LDL uptake by the patients' HMDM to its uptake by control HMDM. Ox-LDL uptake by the patients' HMDM was significantly increased, by 37%, as compared to controls' HMDM (Fig. 4C). PJ supplementation by the patients resulted in a substantial reduction, by 39%, in the uptake of Ox-LDL by the patients' HMDM, as compared to the values obtained before PJ consumption (Fig. 4C).




Fig. 3. The effect of PJ on serum oxidative status and on paraoxonase activity: *in vitro* study. Serum from diabetic patients ( $n = 3$ ) was incubated for 1 h without (-) or with PJ 20  $\mu\text{M}$  or 40  $\mu\text{M}$  polyphenols. Then, the serum was diluted  $\times 4$  with PBS and (A) incubated for 2 h at 37  $^{\circ}\text{C}$  with no addition (basal) or (B) with 100 mM of the free radical generator AAPH (AAPH-induced). At the end of the incubation period, the amount of TBARS in all the samples was determined. (C) The serum from the diabetic patients was incubated with increasing (0–40  $\mu\text{M}$ ) PJ polyphenols concentrations for 1 h at room temperature. Then, paraoxonase 1 arylesterase activity was measured as described under Section 2. Results are given as mean  $\pm$  S.E.M., \* $p < 0.01$  vs. (-); # $p < 0.01$  vs. 0 concentration.

### 3.5. Direct effect of PJ on diabetic patient HMDM oxidative status and on Ox-LDL uptake: *in vitro* study

Incubation of diabetic patient HMDM with PJ (75  $\mu\text{M}$  polyphenols) for 20 h at 37  $^{\circ}\text{C}$  resulted in a significant reduction, by 60%, in the level of cellular peroxides, as measured by the DCFH assay (Fig. 5A). Similarly, incubation of the cells with 75  $\mu\text{M}$  of the PJ-derived polyphenols fraction, significantly decreased the cellular peroxides content in the patients' HMDM by 47% (Fig. 5A). Furthermore, PJ (75  $\mu\text{M}$ ) incubation with diabetic HMDM, inhibited also the uptake of




Fig. 4. The effect of PJ consumption by diabetic patients on human monocytes-derived macrophages (HMDM) oxidative status. Monocytes were isolated from the blood of two healthy subjects (controls) and from three NIDDM patients before and after 3 months of PJ consumption (50 ml per day), as described under Section 2. The monocytes were differentiated into macrophages in the presence of RPMI medium containing 10% autologous serum. After 7 days in culture the amount of cellular peroxides (A) of reduced glutathione (B) and the uptake of Ox-LDL (20  $\mu\text{g}$  of protein/ml) labeled with FITC by the cells (C) were determined as described under Section 2. Results are given as mean  $\pm$  S.E.M., \* $p < 0.01$  vs. controls HMDM.

Ox-LDL by 30%, compared to the extent of Ox-LDL uptake by non-treated cells (Fig. 5B).

## 4. Discussion

The present study demonstrated that pomegranate juice consumption by diabetic patients (as previously shown for healthy subjects and atherosclerotic patients) did not worsen the diabetic parameters, but rather resulted in anti-atherogenic effects with a significant reduction in oxidative stress in the patients' serum and monocytes-macrophages, as well as in macrophage uptake of Ox-LDL.




Fig. 5. The effect of PJ on diabetic patients' monocytes-derived macrophages (HMDM) oxidative status, and on the ability of the cells to take up Ox-LDL: *in vitro* study. Monocytes were isolated from the blood of three NIDDM patients before PJ consumption. After their differentiation into macrophages in the presence of RPMI medium + 10% autologous serum, at day 6, the cells were incubated with no addition (control), with PJ 75  $\mu$ M polyphenols concentration (+PJ), or with 75  $\mu$ M of PJ polyphenols fraction. (A) The level of cellular peroxides was determined by the DCFH assay. (B) The uptake of Ox-LDL (20  $\mu$ g of protein/ml) labeled with FITC was analyzed as described under Section 2. Results are given as mean  $\pm$  S.E.M., \*  $p < 0.01$  vs. control.

Diabetic patients versus control healthy subjects have significant high serum triglyceride levels and low HDL-cholesterol levels, as previously shown [35]. Diabetic patients usually avoid sugar-containing juices which worsen their diabetic markers and atherosclerotic complications. In the present study, we showed that PJ (which contain 10% total sugars [36]) consumption by the diabetic patients did neither increase serum glucose, nor blood HbA1c levels even though PJ glycemic index is similar to that of other fruit juices.

In the patients' serum versus controls we observed increased oxidative stress (high levels of lipid peroxides and aldehydes, and low levels of SH groups), as was previously shown for diabetes [5,37]. The increased serum oxidative stress in the patients could be the result of glycation and glycooxidation of LDL by glucose [38], or/and the decreased capability of the patients' HDL to protect LDL against oxidation [39]. HDL-associated paraoxonase protects LDL and HDL against oxidation, by its ability to hydrolyze specific oxidized lipids [32]. Serum paraoxonase 1 (PON1) activity in our patients versus controls was significantly lower

as was previously shown [40], and this fact may account for the increased serum oxidative stress. Indeed, it was demonstrated that Ox-LDL levels and vascular complications in type II patients are in correlation with PON1 activity [41]. The reduced PON1 activity in the patients could result from PON1 inactivation by oxidized lipids [42], or by glucose [43]. As oxidative stress may have a role in the onset and progression of diabetes and its complications [4–7], anti-oxidants were suggested as a possible treatment [7]. Anti-oxidants such as Vitamin E and red wine were shown to reduce oxidative stress in the patients' serum [8,9]. In accordance with these studies, the present study clearly demonstrated that PJ consumption by diabetic patients significantly reduced serum oxidative stress as shown previously in healthy subjects [13] and in patients with carotid artery stenosis [14]. This effect of PJ may be related to its potent tannins and anthocyanins which scavenge wide spectrum of free radicals [11,12], as well as to PJ-induced increment in the diabetic patients serum PON1 activity as previously shown [13,14]. The increased PON1 activity could have resulted from the reduction in oxidative stress (less PON1 inactivation), and/or from the direct effect of the pomegranate juice component(s) on the enzyme activity as shown *in vitro* (Fig. 3).

Increased oxidative stress was shown not only in the patients' serum, but also in the patients' monocytes–macrophages (increased cellular levels of peroxides and decreased levels of macrophage reduced glutathione). Similar results were noted in peritoneal macrophages from streptozotocin-induced diabetic mice, and also *in vitro* in macrophages that were incubated with high glucose levels [21]. The increased oxidative stress in the patients' HMDM could be related to cellular mechanisms induced by the high glucose levels such as, PKC activation followed by the production of free radicals [3,6,18]. In the present study, PJ consumption significantly reduced cellular peroxides in the patients' HMDM, as was previously shown in carotid lesions from carotid artery stenosis (CAS) patients that consumed PJ [14]. This effect could be due to the increased serum PON1 activity, which can hydrolyze lipid peroxides on the macrophage surface [34,44], or to a direct effect of pomegranate juice component(s). Indeed, *in vitro* we observed that both PJ and the PJ-derived polyphenolic fraction, significantly reduced macrophage oxidative stress. The PJ polyphenolic fraction was less potent than PJ, indicating that other factors in the juice (unique sugars) contribute to cellular oxidative stress reduction. We have recently shown indeed that PJ reduced oxidative stress in J774A.1 macrophage cell line, and this effect was PJ polyphenols dose-dependent [45].

As glucose increases oxidative stress, it was expected that PJ sugars will add to the already increased oxidative stress present in diabetic patients. Surprisingly it did not, and in fact PJ significantly decreased oxidative stress in serum, as well as in their monocytes–macrophages.

The uptake of Ox-LDL by the patients' HMDM was significantly increased as compared to control HMDM. This

phenomenon could be related to the increased expression of the scavenger receptor CD36, which is induced by glucose and/or by the high oxidative stress [34,46]. Similar results were observed in diabetic, streptozotocin-injected mice [21]. PJ consumption by the patients, as well as a direct *in vitro* incubation of PJ with the patients' HMDM resulted in a significant reduction in Ox-LDL uptake by the patients' HMDM. Similar results were noted upon incubating J774A.1 macrophages with PJ [45]. PJ-induced reduction in the cellular uptake of Ox-LDL could not be related to down-regulation of the scavenger receptor CD36 mRNA expression [45]. However, CD36 is not the only surface binding for Ox-LDL by macrophages. PJ component(s) also interact with other scavengers receptors such as the SR-A or LOX-1 (which are also up-regulated by glucose [20,19]), or with proteoglycans which were shown to mediate uptake of Ox-LDL by macrophages [47]. PJ polyphenols could possibly also interfere with the uptake of Ox-LDL by interaction with macrophage surface phospholipids and/or kinases [48].

In conclusion then, pomegranate juice consumption by diabetic patients does not worsen diabetic parameters, but rather act as an anti-atherogenic agent. This anti-atherogenicity is manifested by PJ anti-oxidant properties in serum and monocytes–macrophages, two major components of macrophage foam cell formation, the hallmark of early atherosclerosis.

## Acknowledgment

This study was supported by a grant from the Russell Berrie Foundation, and D-Cure, Diabetes Care in Israel.

## References

- [1] Ziegler D. Type 2 diabetes as an inflammatory cardiovascular disorder. *Curr Mol Med* 2005;5:309–22.
- [2] Wattanakit K, Folsom AR, Selvin E, et al. Risk factors for peripheral arterial disease incidence in persons with diabetes: the Atherosclerosis Risk in Communities (ARIC) Study. *Atherosclerosis* 2005;180:389–97.
- [3] Aronson D, Rayfield EJ. How hyperglycemia promotes atherosclerosis: molecular mechanisms. *Cardiovasc Diabetol* 2002;1:1.
- [4] Ceriello A, Motz E. Is oxidative stress the pathogenic mechanism underlying insulin resistance, diabetes, and cardiovascular disease? The common soil hypothesis revisited. *Arterioscler Thromb Vasc Biol* 2004;24:816–23.
- [5] Martin-Gallan P, Carrascosa A, Gussinye M, Dominguez C. Biomarkers of diabetes-associated oxidative stress and antioxidant status in young diabetic patients with or without subclinical complications. *Free Radic Biol Med* 2003;34:1563–74.
- [6] Whiteside CL. Cellular mechanisms and treatment of diabetes vascular complications converge on reactive oxygen species. *Curr Hypertens Rep* 2005;7:148–54.
- [7] Johansen JS, Harris AK, Rychly DJ, Ergul A. Oxidative stress and the use of antioxidants in diabetes: linking basic science to clinical practice. *Cardiovasc Diabetol* 2005;4:5.
- [8] Park S, Choi SB. Effects of alpha-tocopherol supplementation and continuous subcutaneous insulin infusion on oxidative stress in Korean patients with type 2 diabetes. *Am J Clin Nutr* 2002;75:728–33.
- [9] Ceriello A, Bortolotti N, Motz E, et al. Red wine protects diabetic patients from metal-induced oxidative stress and thrombosis: a pleasant approach to the prevention of cardiovascular disease in diabetes. *Eur J Clin Invest* 2001;31:322–8.
- [10] Rizvi SI, Zaid MA, Anis R, Mishra N. Protective role of tea catechins against oxidation-induced damage of type 2 diabetic erythrocytes. *Clin Exp Pharmacol Physiol* 2005;32:70–5.
- [11] Gil ML, Tomas-Barberan FA, Hess-Pierce B, Holcroft DM, Kader AA. Antioxidant activity of pomegranate juice and its relationship with phenolics composition and processing. *J Agric Food Chem* 2000;10:4581–9.
- [12] Aviram M. Pomegranate juice as a major source for polyphenolic flavonoids and it is most potent antioxidant against LDL oxidation and atherosclerosis. In: Pasquier C, editor. *Proceedings of the XI Biennial Meeting of the Society for Free Radical Research International*. Paris, France, 16–20 July, 2002. Monduzzi Editore S.P. A-MEDIMOND Inc., pp. 523–28.
- [13] Aviram M, Dornfeld L, Rosenblat M, et al. Pomegranate juice consumption reduces oxidative stress, atherogenic modifications to LDL, and platelet aggregation: studies in humans and in atherosclerotic apolipoprotein E-deficient mice. *Am J Clin Nutr* 2000;71:1062–76.
- [14] Aviram M, Rosenblat M, Gaitini D, et al. Pomegranate juice consumption for 3 years by patients with carotid artery stenosis reduces common carotid intima-media thickness, blood pressure and LDL oxidation. *Clin Nutr* 2004;23:423–33.
- [15] Boyle JJ. Macrophage activation in atherosclerosis: pathogenesis and pharmacology of plaque rupture. *Curr Vasc Pharmacol* 2005;3:63–8.
- [16] Naito T, Oikawa S, Kotake H, Hayasaka K, Toyota T. Effect of glucose concentration on foam cell formation in THP-1 cells. *J Atheroscler Thromb* 2001;8:55–62.
- [17] Dobrian AD, Lazar V, Sinescu C, Mincu D, Simionescu M. Diabetic state induces lipid loading and altered expression and secretion of lipoprotein lipase in human monocytes-derived macrophages. *Atherosclerosis* 2000;153:191–201.
- [18] Ceolotto G, Gallo A, Miola M, et al. Protein kinase C activity is acutely regulated by plasma glucose concentration in human monocytes *in vivo*. *Diabetes* 1999;48:1316–22.
- [19] Ling Li, Tatsuya S, Genevieve R. Glucose enhances human macrophage LOX-1 expression: role for LOX-1 in glucose-induced macrophage foam cell formation. *Circ Res* 2004;94:892–901.
- [20] Fukuhara-Takaki K, Sakai M, Sakamoto Y, Takeya M, Horiuchi S. Expression of class A scavenger receptor is enhanced by high glucose *in vitro* under diabetic conditions *in vivo*: one mechanism for an increased rate of atherosclerosis in diabetes. *J Biol Chem* 2005;280:3355–64.
- [21] Hayek T, Hussein K, Aviram M, et al. Macrophage foam cell formation in streptozotocin-induced diabetic mice: stimulatory effect of glucose. *Atherosclerosis*, in press.
- [22] Gan KN, Smolen A, Eckerson HW, La DU BN. Purification of human serum paraoxonase/arylesterase. Evidence for one esterase catalyzing both activities. *Drug Metab Dispos* 1991;19:100–6.
- [23] Frei B, Stocker R, Ames BN. Antioxidant defenses and lipid peroxidation in human blood plasma. *Proc Natl Acad Sci USA* 1988;85:9748–52.
- [24] Buege JA, Aust SD. Microsomal lipid peroxidation. *Methods Enzymol* 1978;52:302–10.
- [25] El-Saadani M, Esterbauer N, El-Sayed M, Goher M, Nassar AY, Jurgens G. Spectrophotometric assay for lipid peroxides in serum lipoproteins using commercially available reagent. *J Lipid Res* 1989;30:627–30.
- [26] Hu M-L, Dillard CJ, Tappel AI. *In vivo* effects of aurothioglucose and sodium thioglucose on rat tissue sulfhydryl levels and plasma sulfhydryl reactivity. *Agents Actions* 1988;25:132–8.
- [27] LeBel CP, Ischiropoulos H, Bondy SC. Evaluation of the probe 2',7'-dichlorofluorescein as an indicator of reactive oxygen



- species formation and oxidative stress. *Chem Res Toxicol* 1992;5:227–31.
- [28] Bass DA, Parce JW, Dechatelet LR, Szejda P, Seeds MC, Thomas M. Flow cytometric studies of oxidative product formation by neutrophils: a graded response to membrane stimulation. *J Immunol* 1983;130:1910–7.
- [29] Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem* 1951;193:265–75.
- [30] Tietze F. Enzymatic method for quantitative determination of nanogram amounts of total and oxidized glutathione: application to mammalian blood and other tissues. *Anal Biochem* 1969;27:502–22.
- [31] Aviram M. Plasma lipoprotein separation by discontinuous density gradient ultracentrifugation in hyperlipoproteinemic patients. *Biochem Med* 1983;30:111–8.
- [32] Aviram M, Rosenblat M. Paraoxonases 1, 2 and 3, oxidative stress and macrophage foam cells formation during atherosclerosis development. *Free Radic Biol Med* 2004;37:1304–16.
- [33] Sies H. Glutathione and its role in cellular functions. *Free Radic Res* 1999;27:916–21.
- [34] Fuhrman B, Volkova N, Aviram M. Oxidative stress increases the expression of the CD36 scavenger receptor and the cellular uptake of oxidized LDL in macrophage from atherosclerotic mice: protective role of antioxidants and paraoxonase. *Atherosclerosis* 2002;161:307–16.
- [35] Krentz AJ. Lipoprotein abnormalities and their consequences for patients with type 2 diabetes. *Diab Obes Metab* 2003;5(Suppl. 1):S19–27.
- [36] El-Nemr SE, Ismail IA, Ragab M. Chemical composition of juice and seeds of pomegranate fruit. *Nahrung* 1991;34:601–6.
- [37] Ozdemir G, Ozden M, Maral H, Kuskay S, Cetinalp P, Tarkum I. Malondialdehyde, glutathione, glutathione peroxidase and homocysteine levels in type 2 diabetic patients with and without microalbuminuria. *Am Clin Biochem* 2005;42:99–104.
- [38] Knott HM, Brown BE, Davies MJ, Dean RT. Glycation and glycooxidation of low-density lipoproteins by glucose and low-molecular mass aldehydes. Formation of modified and oxidized particles. *Eur J Biochem* 2003;270:3572–82.
- [39] Gowri MS, Van der Westhuyzen DR, Brodges SR, Anderson JW. Decreased protection by HDL from poorly controlled type 2 diabetic subjects against LDL oxidation may be due to the abnormal composition of HDL. *Arterioscler Thromb Vasc Biol* 1999;19:2226–33.
- [40] Letellier C, Duron MR, Jouanolle AM, et al. Serum paraoxonase activity and paraoxonase gene polymorphism in type 2 diabetic patients with or without vascular complications. *Diab Metab* 2002;28:297–304.
- [41] Tsuzura S, Ikeda Y, Suchiro T, et al. Correlation of plasma oxidized low-density lipoprotein levels to vascular complications and human serum paraoxonase in patients with type 2 diabetes. *Metabolism* 2004;53:297–302.
- [42] Aviram M, Rosenblat M, Billecke S, et al. Human serum paraoxonase (PON 1) is inactivated by oxidized low density lipoprotein and preserved by antioxidants. *Free Radic Biol Med* 1999;26:892–904.
- [43] Ferretti G, Bacchetti T, Marchionni C, et al. Effect of glycation of high density lipoproteins on their physicochemical properties and on paraoxonase activity. *Acta Diabetol* 2001;38:163–9.
- [44] Rozenberg O, Shih DM, Aviram M. Paraoxonase (1) attenuates macrophage oxidative status: studies in PON1 transfected cells and in PON1 transgenic mice. *Atherosclerosis* 2005;181:9–18.
- [45] Fuhrman B, Volkova N, Aviram M. Pomegranate juice inhibits oxidized LDL uptake and cholesterol biosynthesis in macrophages. *Atherosclerosis* 2005;16:570–6.
- [46] Sampson MJ, Davies IR, Braschi S, Ivory K, Hughes DA. Increased expression of scavenger receptor (CD36) in monocytes from subjects with type 2 diabetes. *Atherosclerosis* 2003;167:120–34.
- [47] Kaplan M, Williams KJ, Mandel H, Aviram M. Role of macrophage glycosaminoglycans in the cellular catabolism of oxidized LDL by macrophages. *Arterioscler Thromb Vasc Biol* 1998;4:542–53.
- [48] Sah JF, Balasubramaniam S, Ecker RL, Rorke EA. Epigallocatechin-3-gallate inhibits epidermal growth factor receptor signaling pathway. Evidence for direct inhibition of ERK1/2 and AKT kinases. *J Biol Chem* 2004;279:12755–62.